

COMPOSITION LAW OF CARDINAL ORDERING PERMUTATIONS

Jesús San Martín ^{a,b} M^a Jose Moscoso ^a

^a*Departamento de Matemática Aplicada, E.U.I.T.I. Universidad Politécnica de Madrid. 28012-Madrid*

^b*Departamento de Física Matemática y de Fluidos, Facultad de Ciencias. Universidad Nacional de Educación a Distancia. 28040-Madrid, SPAIN.*

A. González Gómez ^c

^c*Departamento de Matemática Aplicada a los Recursos Naturales, E.T. Superior de Ingenieros de Montes. Universidad Politécnica de Madrid. 28040-Madrid, SPAIN.*

Abstract

In this paper the theorems that determine composition laws for both cardinal ordering permutations and their inverses are proven. So, the relative positions of points in a hs -periodic orbit become completely known as well as in which order those points are visited. No matter how a hs -periodic orbit emerges, be it through a period doubling cascade ($s = 2^n$) of the h -periodic orbit, or as a primary window (like the saddle-node bifurcation cascade with $h = 2^n$), or as a secondary window (the birth of a s -periodic window inside the h -periodic one). Certainly, period doubling cascade orbits are particular cases with $h = 2$ and $s = 2^n$. Both composition laws are also shown in algorithmic way for their easy use.

Key words: Cardinal ordering permutations. Composition law. Preceding sequence.

1 Introduction

As Hao Bai-Lin says in his book [1] “Symbolic dynamics might be the only rigorous way to define chaos...”. Certainly, this nice theory brings order into

Email addresses: `jsm@dfmf.uned.es` (Jesús San Martín), `mariajose.moscoso@upm.es` (M^a Jose Moscoso), `antonia.gonzalez@upm.es` (A. González Gómez).

chaos labelling and ordering periodic orbits and therefore giving us the grammar of chaos. Working in this area, Metropolis, Stein and Stein proved, in their seminal work [2], that the transformations of an interval in itself, by an unimodal function f , generate limit sets exhibiting universal structures. The limit sets are characterized by finite sequences of symbols R and L, that indicate whether successive iterates of the unimodal function are plotted right or left of the critical point of the function. If a sequence of symbols is repeated periodically then it represents a periodic orbit of the dynamical system.

Once symbolic sequences associated to orbits are known a new question arises. How are symbolic sequences of h -periodic and s -periodic orbits composed to obtain the symbolic sequence of a hs -periodic orbit?. The answer to this problem is found in the fundamental work by Derrida, Gervois and Pomeau [3]. The Derrida-Gervois-Pomeau composition law allows us to get new information about a dynamical system from information already known. It is an example of the power of symbolic dynamics, that we want to generalize in this paper.

If we pay attention to symbolic sequence of a h -periodic orbit we observe a sequence of letters R and L, new questions, logically, arise: what are the relative positions of different R(L)? what R(L) is rightmost or leftmost than other R(L)? is a given R(L) to the right or left of another previously given R(L)?. To answer these questions we need to introduce an order on the set of points that form the h -periodic orbit. This order determines the relative positions of points about which questions are answered. This problem was approached and solved in [4] for period doubling cascade [5] [6] (see kneading theory [7] and invariant coordinates [8]) for a different approach). In this same paper, the authors met the cardinal ordering permutation σ_{2^k} for every orbit of period doubling cascade. This permutation allows to know both how many iterates are needed to reach a given point of the orbit from another fixed point and the relative position of the point.

Bearing in mind that the symbolic sequence composition law provides information about new orbits then the natural generalization is how to compose two arbitrary cardinal ordering permutation σ_h and σ_s to obtain σ_{hs} . From this generalization visiting order and relative position of points in a hs -periodic orbit are obtained, that is, new information about the dynamical system is deduced. In fact, as it is discussed in the conclusions, infinitely many orbits are fully determined by only two cardinal ordering permutations. Furthermore, cardinal ordering permutations for period doubling cascade and saddle-node bifurcation cascade [9,10] are immediately derived and therefore all underlying periodic structures in an unimodal map. That is, these structures are fully determined from their cardinal ordering permutations that we can obtain with the composition law. The most remarkable goal of this paper is to find such a composition law.

Before starting the main part of the paper we are going to introduce some intuitive ideas and related problems that will be fundamental in its development and useful to understand the sometimes tedious mathematical apparatus.

If the hs -periodic orbit is the result of the birth of a s -periodic orbit in a h -periodic window, there is an intuitive and geometric understanding of the problem that we will use to obtain the composition law. Imagine there are h rooms in a corridor, bearing s chairs each. Every time you move an iterate you go into a different room and you sit in a chair in the room. You do not visit the rooms one after the another, as found in the corridor, but according to a rule given by σ_h . Each time you visit the same room, you will sit in a different chair according to σ_s .

Two questions naturally arise:

- (1) How many movements (iterates) do we have to make to reach the r -th chair (supposed they were arranged in a row)?
- (2) What chair would you reach if you made q movements (iterates)?

Both questions will be answered in this paper. They are very related to composition law. That can give an idea of the power of the law. Nonetheless to get the answers two important problems have to be previously solved.

- i) There are two kinds of rooms, associated with a maximum and minimum of f^h respectively (every iteration of f represents a movement). When we enter in a room of maximum kind then σ_s indicates on what chair we sit down, but if we enter in a room of minimum kind we must use the conjugate of the permutation $\overline{\sigma_s}$ [4]. Therefore, we have to know whether f^h has a maximum or a minimum. Should f^h had 5000 extrema and we had to take the 5000 derivatives of f^h the method would be totally useless. We have to get the information easily and without taking derivatives, so we will prove a proposition that allows us to get this information very easily.
- ii) If we are in a room seated in a chair then we know what chair we will sit the next time we visit the room, because of σ_s . But, if we visit the room for the first time we do not have that information. Therefore, it is necessary to determine what chair we will sit the first time we visit a room.

The solutions of both problems are related and they are an important milestone in finding the composition law of cardinal ordering permutation.

Apart from its theoretical value *per se*, a composition law for permutations is useful wherever permutation ruling orbits are used, since composition law indicates how to compound different elements to get more information. Its properties are expected to underlie in permutations modelling area preserving monotone twist maps [11], entropy via permutations [12,13,14,15] and patterns in dynamical system [16,17]; subjects, that in turn, underlie in other fields [18].

This paper is organized as follows. First, most of the definitions are introduced. Second, some lemmas and proposition are proven to solve the mentioned problems. Thirst, composition laws for cardinal ordering permutations are obtained. Then composition laws are shown in algorithmic way. Finally the last section contains our conclusions and discussions. Some examples will also be shown to ease the use of theorems and algorithms by scientists and engineers.

2 Definitions and notation

Let $f : I \subset \mathbb{R} \rightarrow I$ be an unimodal map, and let C denote its critical point (we suppose, without loss of generality, that C is a maximum). Let $O_p = \{x_1, \dots, x_p \equiv C\} = \{f(C), \dots, f^{p-1}(C), f^p(C) \equiv C\}$ be a p -periodic supercycle of f , then the first and the second iterates of C determine the subinterval $J = [f^2(C), f(C)]$ such that $O_p \subset [f^2(C), f(C)] = J$. We denote $J_L = [f^2(C), C]$ and $J_R = [C, f(C)]$. Since the orbit O_p does not have the natural order within the interval $J = [f^2(C), f(C)]$, some previous definitions, introduced in [4], are needed.

The set $\{C_{(1,p)}^*, C_{(2,p)}^*, \dots, C_{(p,p)}^*\}$ will denote the descending cardinality ordering of the orbit $O_p = \{x_1, x_2, \dots, x_p \equiv C\}$. Given that $C_{(i,p)}^*$ $i = 1, \dots, p$, is the point of the orbit O_p located in the position ' i ' according to the descending cardinality ordering, the point $C_{(i,p)}^*$ is defined as the i -th cardinal of the p -period orbit. Notice that $f^2(C) = C_{(p,p)}^* < \dots < C_{(2,p)}^* < C_{(1,p)}^* = f(C)$.

Let O_p be a p -periodic supercycle of f , we denote as σ_p the permutation $\sigma_p = (\sigma(1, p), \dots, \sigma(p, p))$, where $\sigma(i, p)$ is the number of iterations of f from C such that $f^{\sigma(i,p)}(C) = C_{(i,p)}^* = x_{\sigma(i,p)}$ for all $i = 1, \dots, p$ (see figures 1 and 2). This permutation will be called the cardinal ordering permutation of the orbit O_p .

Authors proved in [4] that the inverse permutation determines the order in which the orbit is visited. That is, the visiting sequence of the p -periodic supercycle O_p from C is the inverse permutation $\sigma_p^{-1} = (\sigma^{-1}(1, p), \sigma^{-1}(2, p), \dots, \sigma^{-1}(p, p))$ (see figures 1 and 2). Therefore, depending on which context it is used, σ_p^{-1} will be called inverse permutation or visiting sequence.

Definition 1 *The symbolic sequence of the supercycle O_p , denoted by $I_1, I_2, \dots, I_{p-1}, C$ where $I_i = R$ or L for all $i = 1, \dots, p-1$, will be called p -symbolic sequence.*

Definition 2 *Let $I_1, I_2, \dots, I_{p-1}, C$ be a p -symbolic sequence. Then*

(i) I_j is the p -symbolic sequence element associated to the element x_j of the




Figure 1. $\sigma_3 = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}$ $\leftarrow$ indicates the position ' i ' of $C_{(i,3)}^*$ $\leftarrow$ number of iterations, $\sigma_{(i,3)}$, to reach $C_{(i,3)}^*$ $\sigma_3^{-1} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}$




Figure 2. $\sigma_4 = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 4 & 3 & 2 \end{pmatrix}$ $\sigma_4^{-1} = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 4 & 3 & 2 \end{pmatrix}$




Figure 3. The graph of f^s is repeated close to the straight line $y = x$

orbit O_p .

- (ii) $I_1, I_2, \dots, I_{j-1}$ is the subsequence preceding I_j of the p -symbolic sequence. The subsequence preceding I_1 of the p -symbolic sequence is C by definition.
- (iii) $I_j, \dots, I_{p-1}, C$ is the subsequence following I_{j-1} of the p -symbolic sequence.
- (iv) $I_j, I_{j+1}, \dots, I_{q-1}, I_q$ is the p -symbolic sequence from I_j to I_q .

We want to study a hs -periodic supercycle O_{hs} of f with hs points, resulting from the combination of two supercycles with h and s points respectively. In order to get that we split the hs points in h sets, that we will name boxes, with s points each. The boxes are visited according to σ_h^{-1} , while σ_s^{-1} gives the order in which points inside the boxes are visited. As the boxes are visited one after another according σ_h^{-1} , it results that the $h-1$ first elements of the symbolic sequence of O_{hs} coincide with the elements of the symbolic sequence of O_h . The boxes are associated with the extrema of f^h (closest to the straight line $y = x$) and each of them contains the extrema of f^s (closest to the straight line $y = x$) as it can be seen in figure 3. Notice that the graph of f^s is repeated in each extremum of f^h , and furthermore they intersect the straight line $y = x$: the points of the supercycle O_{hs} .

According to this idea we introduce the following definition

Definition 3 Let $f(x; r)$ be an unimodal map such that for $r = r_h$ and $r = r_s$ it has the supercycles O_h and O_s having periods h and s , respectively. We define hs -periodic supercycle of f , denoted by O_{hs} , the supercycle of f $O_{hs} =$

$\{x_1, x_2, \dots, x_{hs} \equiv C\}$ originated at $r = r_{hs}$, where:

- i) The hs points split in h sets with s points each.
- ii) The h sets are in the neighborhoods of the extrema of $f^h(x; r_{hs})$, closest to the straight line $y = x$, containing the graph of $f^s(x; r_s)$.

The definition 3 refers to supercycles associated with period doubling cascade, saddle-node bifurcation cascade and the birth of windows inside other windows.

Definition 4 Let $\{C_{(1,hs)}^*, C_{(2,hs)}^*, \dots, C_{(hs,hs)}^*\}$ be the cardinal points of O_{hs} . The i -th cardinal box of O_{hs} with $i = 1, \dots, h$, denoted by H_i^* , is the set $H_i^* = \{c_{(1,i)}^*, c_{(2,i)}^*, \dots, c_{(s,i)}^*\}$, where

$$c_{(j,i)}^* = C_{((i-1)s+j,hs)}^* \quad \forall j = 1, \dots, s$$

Observe that elements belonging O_{hs} are denoted by capital letters, but when these elements belong to a box they are denoted by small letters. With this definition we achieve that both the boxes H_i^* with $i = 1, \dots, h$ and the points inside them maintain the descending cardinality ordering.

Remark 1 As said above σ_h^{-1} determines how boxes are visited, therefore if $x_j \in O_{hs}$ with $1 \leq j \leq h$ then $x_j \in H_{\sigma^{-1}(j,h)}^*$. In fact, x_{j+rh} with $r = 0, \dots, s-1$ are the unique points belonging to $H_{\sigma^{-1}(j,h)}^*$. In other words we can say that if $x_j \in H_i^*$ then $j = \sigma(i, h) + rh$ for some $r = 0, \dots, s-1$.

Definition 5 The central box of O_{hs} , denoted as H_C^* , is the box containing C , that is $H_C^* = H_{\sigma_{(h,h)}^{-1}}^*$. Notice that, according to remark 1, $x_{rh} \in H_C^* = H_{\sigma_{(h,h)}^{-1}}^*$ with $r = 1, \dots, s$.

We must point out that $H_{\sigma_{(j,h)}^{-1}}^* \neq H_C^*$ with $1 \leq j < h$, then either $H_{\sigma_{(j,h)}^{-1}}^* \subset J_L$ or $H_{\sigma_{(j,h)}^{-1}}^* \subset J_R$. This result will be very useful and widely used along the paper.

Definition 6 The box H_i^* is a convex box of O_{hs} if there exists some $x_k \in H_i^* \subset O_{hs}$ such that x_k is a maximum of f^h . The box H_i^* is a concave box of O_{hs} if there exists some $x_k \in H_i^* \subset O_{hs}$ such that x_k is a minimum of f^h .

Definition 7 We say that $c_{(1,i)}^*$ and $c_{(s,i)}^*$ are boundary points for all $i = 1, \dots, h$.

Definition 8 Let be $O_{hs} = \{x_1, x_2, \dots, x_{hs} \equiv C\}$. We say that the order entry from $H_{\sigma^{-1}(n,h)}^*$ to $H_{\sigma^{-1}(m,h)}^*$, with $1 \leq n < m \leq h$, is conserved if $f^{m-n}(c_{(1,\sigma^{-1}(n,h))}^*) = c_{(1,\sigma^{-1}(m,h))}^*$ or $f^{m-n}(c_{(s,\sigma^{-1}(n,h))}^*) = c_{(s,\sigma^{-1}(m,h))}^*$.

We say that the order entry from $H_{\sigma^{-1}(n,h)}^*$ to $H_{\sigma^{-1}(m,h)}^*$, with $1 \leq n < m \leq h$,

is reversed if $f^{m-n}(c_{(1,\sigma^{-1}(n,h))}^*) = c_{(s,\sigma^{-1}(m,h))}^*$ or $f^{m-n}(c_{(s,\sigma^{-1}(n,h))}^*) = c_{(1,\sigma^{-1}(m,h))}^*$.

3 Theorems

In order to get our goals we need to prove some lemmas that make easy the proof of the theorems.

Lemma 1 *Let $O_{hs} = \{x_1, x_2, \dots, x_{hs} \equiv C\}$ be a hs -periodic supercycle of f . Then $x_1, x_2, \dots, x_h$ are boundary points.*

Proof 1 *As $x_1 = C_{(1,hs)}^* = c_{(1,1)}^*$ and $x_2 = C_{(hs,hs)}^* = c_{(s,h)}^*$ it results that x_1 and x_2 are boundary points from definition 7.*

Let us first prove that x_3 is also a boundary points. We have $x_2 = c_{(s,h)}^$ then $x_2 \in H_h^* = H_{\sigma^{-1}(2,h)}^*$ thereby $x_2 < x_{2+rh}$ for all $r = 1, \dots, s-1$. Given $H_h^* \subset J_L$ and f is an increasing function in J_L it results that $x_3 = f(x_2) < f(x_{2+rh})$. As σ_h^{-1} determines how boxes are visited we obtain that $x_3, f(x_{2+rh}) \in H_{\sigma^{-1}(3,h)}^*$ and as $x_3 < f(x_{2+rh})$ it yields $x_3 = c_{(s,\sigma^{-1}(3,h))}^*$ and from definition 7 it results x_3 is a boundary point.*

Let us next prove that x_4 is also a boundary point. As $x_3 = c_{(s,\sigma^{-1}(3,h))}^ \in H_{\sigma^{-1}(3,h)}^*$ then we have $x_3 < x_{3+rh}$ for all $r = 1, \dots, s-1$. Two cases are possible:*

i) $H_{\sigma^{-1}(3,h)}^* \subset J_L$.

In this case we repeat the previous argument and obtain $x_4 = c_{(s,\sigma^{-1}(4,h))}^$.*

ii) $H_{\sigma^{-1}(3,h)}^* \subset J_R$.

As f is a decreasing function in J_R it results $f(x_3) > f(x_{3+rh})$ for all $r = 1, \dots, s-1$. Since σ_h^{-1} determines how boxes are visited and furthermore $f(x_3) = x_4, f(x_{3+rh}) \in H_{\sigma^{-1}(4,h)}^$ then it yields that $x_4 = c_{(1,\sigma^{-1}(4,h))}^*$ and, from definition 7, x_4 is a boundary point.*

The same proof works for $x_5, \dots, x_h$.

Example 1 *If we pay attention to figures 4 and 5, where a $O_{3,4}$ supercycle is plotted, we can see that $x_1 = f(C)$, $x_2 = f(x_1)$ and $x_3 = f(x_2)$ are boundary points. They are spotted in the boundaries of their respective boxes.*

In order to prove the next lemma we want to point out that if $1 \leq n < m \leq h$, $n, m \in \mathbb{N}$, then by the above lemma we will only work with boundary points; hence, “entry order is not conserved” is equivalent to “entry order is reversed”.

Lemma 2 *Assuming $1 \leq n < m \leq h$, $n, m, h \in \mathbb{N}$, the entry order from $H_{\sigma^{-1}(n,h)}^*$ to $H_{\sigma^{-1}(m,h)}^*$ is conserved if only if the R-parity of the hs -symbolic*




Figure 4. An $O_{3,4}$ supercycle can be observed. Boxes H_1^* , H_2^* , H_3^* have been plotted in the extrema of f^3 . Some cardinals have also been plotted. $C_{5,12}^*$ and $C_{8,12}^*$ are boundary points in H_2^* .

sequence from I_n to I_{m-1} is even.

Proof 2 Suppose the R-parity of the hs -symbolic sequence from I_n to I_{m-1} is even.

Let $x_n \in H_{\sigma^{-1}(n,h)}^*$ with $1 \leq n < m \leq h$. According to lemma 1, x_n is a boundary point. Without loss of generality, let $x_n = c_{(1,\sigma^{-1}(n,h))}^*$ and I_n the hs -symbolic sequence element associated to x_n .

It is deduced from proof of lemma 1 that

$$f(c_{(1,\sigma^{-1}(n,h))}^*) = \begin{cases} c_{(1,\sigma^{-1}(n+1,h))}^* & \text{if } H_{\sigma^{-1}(n,h)}^* \subset J_L \\ c_{(s,\sigma^{-1}(n+1,h))}^* & \text{if } H_{\sigma^{-1}(n,h)}^* \subset J_R \end{cases}$$




Figure 5. Enlargement of the box H_1^* in figure 4, where boundary points $C_{1,12}^*$ and $C_{4,12}^*$ can be seen

$$f(c_{(s,\sigma^{-1}(n,h))}^*) = \begin{cases} c_{(s,\sigma^{-1}(n+1,h))}^* & \text{if } H_{\sigma^{-1}(n,h)}^* \subset J_L \\ c_{(1,\sigma^{-1}(n+1,h))}^* & \text{if } H_{\sigma^{-1}(n,h)}^* \subset J_R \end{cases}$$

As $I_n = L$ if and only if $H_{\sigma^{-1}(n,h)}^* \subset J_L$ and $I_n = R$ if and only if $H_{\sigma^{-1}(n,h)}^* \subset J_R$ (meanwhile $H_{\sigma^{-1}(n,h)}^*$ is not the central box because the central box is the last visited one), we have that

$$f(c_{(1,\sigma^{-1}(n,h))}^*) = \begin{cases} c_{(1,\sigma^{-1}(n+1,h))}^* & \text{if } I_n = L \\ c_{(s,\sigma^{-1}(n+1,h))}^* & \text{if } I_n = R \end{cases}$$

$$f(c_{(s,\sigma^{-1}(n,h))}^*) = \begin{cases} c_{(s,\sigma^{-1}(n+1,h))}^* & \text{if } I_n = L \\ c_{(1,\sigma^{-1}(n+1,h))}^* & \text{if } I_n = R \end{cases}$$

Thus, if $I_n = L$ the entry order from $H_{\sigma^{-1}(n,h)}^*$ to $H_{\sigma^{-1}(n+1,h)}^*$ is conserved; whereas that if $I_n = R$ the entry order is reversed.

Using the previous argument it successively results that if the R -parity of the hs -symbolic sequence from I_n to I_{m-1} is even then $f^{m-n}(c_{(1,\sigma^{-1}(n,h))}^*) =$

$c_{(1,\sigma^{-1}(m,h))}^*$, that is, the entry order from $H_{\sigma^{-1}(n,h)}^*$ to $H_{\sigma^{-1}(m,h)}^*$ is conserved and necessary condition has been proved.

On the other hand if the R-parity of the hs -symbolic sequence from I_n to I_{m-1} is odd then, by using the above argument it is concluded that $f^{m-n}(c_{(1,\sigma^{-1}(n,h))}^*) = c_{(s,\sigma^{-1}(m,h))}^*$, that is, the entry order from $H_{\sigma^{-1}(n,h)}^*$ to $H_{\sigma^{-1}(m,h)}^*$ is reversed. Note that we have actually proved the sufficiency condition.

Example 2 Let us pay attention to figure 4. The R-parity from I_1 (associated to $x_1 = C_{(1,12)}^*$) to I_2 (associated to $x_2 = C_{(12,12)}^*$) is odd therefore according to lemma 2 the entry order is reversed, as we can see in figure 4, paying attention to boxes H_1 and H_3 . The R-parity from I_2 (associated to $x_2 = C_{(12,12)}^*$) to I_3 (associated to $x_3 = C_{(8,12)}^*$) is even therefore, according to lemma 2, the entry order is conserved, as we can see in figure 4, paying attention to boxes H_3 and H_2 .

The lemma, we have just proven, allows us to know what point in a box is reached the first time that box is visited. The following times we visit that box we will know what point is reached by using σ_s^{-1} or $\overline{\sigma_s^{-1}}$ depending on whether the box is associated with a maximum or a minimum of f^h . This kind of information will be obtained in lemma 3 for which we need to introduce the following definition.

Definition 9 Let $I_1, \dots, I_{p-1}, C$ be a p -symbolic sequence. We define the sign of p -symbolic sequence, and we denote as $\text{sgn}(I_1, \dots, I_{p-1}, C)$, as the number given by

$$\text{sgn}(I_1, \dots, I_{p-1}, C) = \begin{cases} 1 & \text{if R-parity of } I_1, \dots, I_{p-1} \text{ is even} \\ -1 & \text{otherwise} \end{cases}$$

Lemma 3 Let $O_p = \{x_1, \dots, x_p \equiv C\}$ be a p -periodic supercycle of an unimodal function f with a maximum at C . Then f^p has a maximum (minimum) at x_i , for all $i = 1, \dots, p$, if and only if the R-parity of the subsequence preceding I_i in the p -symbolic sequence is even (odd).

Proof 3 As we are working with a supercycle, then $(f^p(x_i))' = 0$ for all $i = 1, \dots, p$ and all the points belonging to the supercycle are either maxima or minima. Thus, we only need to study the sign of $(f^p(x_i))''$ at each $x_i \in O_p$ to know whether we have a maximum or a minimum.

By using the chain rule and taking into account that $(f^{p-i})'(x_i) = f'(C) = 0$ it yields:

$$(f^p(x_i))'' = f''(f^{p-i}(x_i)) \prod_{j=1}^{i-1} f'(f^{p-j}(x_i)) \prod_{j=i+1}^p (f'(f^{p-j}(x_i)))^2$$

As f has a maximum at C it follows $f''(f^{p-i}(x_i)) = f''(C) < 0$ and it implies

$$\text{sgn}((f^p(x_i))'') = -\text{sgn}\left(\prod_{j=1}^{i-1} f'(f^{p-j}(x_i))\right)$$

consequently,

$$\text{sgn}((f^p(x_i))'') = -\text{sgn}\left(\prod_{j=1}^{i-1} f'(x_j)\right)$$

Since $f'(x_i) > 0$ if $I_i = L$ and $f'(x_i) < 0$, if $I_i = R$, then

$$\text{sgn}\left(\prod_{j=1}^{i-1} f'(x_j)\right) = \text{sgn}(I_1, \dots, I_{i-1}, C)$$

thus, we obtain

$$\text{sgn}((f^p(x_i))'') = -\text{sgn}(I_1, \dots, I_{i-1}, C) \quad (1)$$

So if f^p has a maximum at x_i then $\text{sgn}((f^p(x_i))'') = -1$. Therefore, from (1) we obtain

$$-1 = -\text{sgn}(I_1, \dots, I_{i-1}, C)$$

Consequently, the R -parity of the subsequence preceding I_i is even.

On the other hand, if the R -parity of the subsequence preceding I_i is even then from (1) we have

$$\text{sgn}((f^p(x_i))'') = -1$$

and f^p has a maximum at x_i .

Notice that we can analogously prove that f^p has a minimum at x_i if and only if the R -parity of the subsequence preceding I_i is odd.

Example 3 In figure 1 can be seen a 3-periodic supercycle with symbolic sequence RLC . The R -parity of the subsequence preceding x_1 is even, and f^3 has a maximum at this point. Meanwhile the R -parity of the subsequences preceding x_2 and x_3 are odd and f^3 has minima at these points.

Lemma 3 allows us to know whether a box is a concave or a convex one, without deriving f^h . We need to know this information because depending

on it σ_s^{-1} or $\overline{\sigma_s^{-1}}$ is used to jump from one point to another inside a box. Obviously it takes less time to calculate a previous sequence that deriving f^h twice, but if h is big then calculating a preceding subsequence can take a long time too. However former lemmas can be exploit to get a more refined information as the below proposition means:

“A box is convex (concave) if and only if the first time the box is visited a $c_{(1,-)}^*$ ($c_{(s,-)}^*$) point is reached”. That is, the box is convex or concave depending on whether the first time you visit the box you reach the first or the last boundary point. That in itself hides an unexpected link between analysis and topology in dynamical systems that it must be researched.

Before proving the proposition we must take into account the following remark.

Remark 2 *Let $O_{hs} = \{x_1, x_2, \dots, x_{hs} \equiv C\}$ be a hs -periodic supercycle of f , then $(f^{hs})'(x_i) = 0$ for each $i = 1, \dots, hs$. As f is an unimodal function with an unique critical point at C , it is deduced that there exists $\{x'_1, x'_2, \dots, x'_h \equiv C\} \subset O_{hs}$ such that $(f^h)'(x'_j) = 0$ for each $j = 1, \dots, h$. These points determine the boxes according to definition 6.*

If we chose any point belonging to the central box after $h - 1$ iterates it would generate a symbolic sequence coinciding with the $h - 1$ first letters of the symbolic sequence of O_h , that also coincides with the $h - 1$ first letters of the symbolic sequence of O_{hs} . That is so because each box of O_{hs} is visited according to σ_h^{-1} , the permutation that determines the supercycle O_h and its symbolic sequence.

Proposition 4 *Let $O_{hs} = \{x_1, x_2, \dots, x_{hs} \equiv C\}$ be a hs -periodic supercycle of f and let $x_r \in H_{\sigma^{-1}(r,h)}^*$ with $1 \leq r \leq h$, then*

- (i) $H_{\sigma^{-1}(r,h)}^*$ is a convex box if and only if $x_r = c_{(1,\sigma^{-1}(r,h))}^*$
- (ii) $H_{\sigma^{-1}(r,h)}^*$ is a concave box if and only if $x_r = c_{(s,\sigma^{-1}(r,h))}^*$

Proof 4 (i) *Let us suppose that $H_{\sigma^{-1}(r,h)}^*$ is a convex box, then there exists some $x_k \in O_{hs}$ with $x_k \in H_{\sigma^{-1}(r,h)}^*$ such that f^h has a maximum at x_k .*

Let $O_h = \{x'_1, \dots, x'_h \equiv C\}$ be the h -periodic supercycle of f , as $x_k \in H_{\sigma^{-1}(r,h)}^$, according to remark 2 it must be $x_k = x'_r$. As there is a maximum at $x_k = x'_r$ it yields, by using lemma 3, that the R -parity of $I'_1, \dots, I'_{r-1}$ is even. Since $I'_1, \dots, I'_{r-1}$ coincides with the first $r - 1$ symbols of the hs -symbolic sequence, that is, $I_1, \dots, I_{r-1}$, then the R -parity of $I_1, \dots, I_{r-1}$ is also even. Therefore, it results, after applying lemma 2, that the entry order from $H_{\sigma^{-1}(1,h)}^*$ to $H_{\sigma^{-1}(r,h)}^*$ is conserved. As $x_1 = c_{(1,1)}^* \in H_{\sigma^{-1}(1,h)}^* = H_1^*$ and the entry order is conserved then it is concluded that $x_r = c_{(1,\sigma^{-1}(r,h))}^*$.*

The inverse implication can be obtained by reversing the process.

(ii) *The proof is exactly the same as before.*

Notice we have never said x'_r is a boundary point and, in fact, it is not. The idea of the proof is that every extremum of f^h (the boxes) behaves as f . The composition (s times) of this approximation generates something similar to f^s (the points inside the boxes) in the neighborhood of the extrema (see figure 3). Then we link an extremum of f^h (at x'_r) with a boundary point of that “thing similar to f^s ” (around x_r). Obviously x'_r and x_r belong to the same box generated by the extremum of f^h at x'_r ; in fact, σ_s^{-1} moves x'_r to x_r (maximum at x'_r) and $\overline{\sigma_s^{-1}}$ moves x'_r to x_r (minimum at x'_r) because x'_r behaves as the critical point C. Therefore instead of getting information from x'_r (behaving as C, an extremum), we get it from displacement of x'_r (a boundary point) given by σ_s^{-1} .

Example 4 *Let us pay attention to figures 4 and 5. H_1^* is a convex box because the first time this box is visited the point $C_{(1,12)}^* = c_{(1,1)}^*$ is reached. Meanwhile H_2^* and H_3^* are concave boxes because the first time these boxes are visited the points $C_{(12,12)}^* = c_{(4,3)}^*$ and $C_{(8,12)}^* = c_{(4,2)}^*$ are respectively reached.*

As we have said, the boxes, associated to extrema of f^h , can be either convex (maximum of f^h) or concave (minimum of f^h). The permutation σ_s^{-1} determines the visiting order in convex boxes. To determine the visiting order in a concave box the permutation σ_s^{-1} must be transformed by the same process that transforms a maximum into a minimum: a homotecy, as it can be seen in Appendix. According to this idea we have the conjugated permutation as shown below.

Definition 10 *Let σ_p be the cardinal ordering permutation of the p -periodic supercycle of f . We define the cardinal ordering conjugated permutation, denoted by $\overline{\sigma_p}$, as*

$$\overline{\sigma_p} = \begin{pmatrix} 1 & \cdots & i & \cdots & p \\ \sigma(p, p) & \cdots & \sigma(p - i + 1, p) & \cdots & \sigma(1, p) \end{pmatrix}$$

Definition 11 *Let σ_p^{-1} be the visiting sequence of the p -periodic supercycle O_p . We define the conjugated visiting sequence, denoted by $\overline{\sigma_p^{-1}}$, as*

$$\overline{\sigma_p^{-1}} = \begin{pmatrix} 1 & \cdots & i & \cdots & p \\ p + 1 - \sigma^{-1}(1, p) & \cdots & p + 1 - \sigma^{-1}(i, p) & \cdots & p + 1 - \sigma^{-1}(p, p) \end{pmatrix}$$

Theorem 5 *Let O_h and O_s be supercycles with cardinal points $\{C_{(1,h)}^*, \dots, C_{(h,h)}^*\}$ and $\{C_{(1,s)}^*, \dots, C_{(s,s)}^*\}$ respectively. Let σ_h^{-1} and σ_s^{-1} be their visiting sequences and let $m \in \mathbb{N}$ where $1 \leq m \leq hs$ and $m = nh + r$, $1 \leq r \leq h$. Then the visiting sequence of O_{hs} is,*

- i) if the R-parity of the subsequence preceding I_r of the h -symbolic sequence is even

$$\sigma^{-1}(m, hs) = (\sigma^{-1}(r, h) - 1)s + \sigma^{-1}(n + 1, s)$$

- ii) if the R-parity of subsequence preceding I_r of the h -symbolic sequence is odd

$$\sigma^{-1}(m, hs) = (\sigma^{-1}(r, h) - 1)s + s + 1 - \sigma^{-1}(n + 1, s)$$

Proof 5 To know the cardinal point that we are located after m iterations from C , we can write $f^m(C) = f^{nh}(f^r(C))$.

As $f^r(C) = x_r \in H_{\sigma^{-1}(r, h)}^*$, using lemma 1, it yields x_r is a boundary point, that is, $x_r = c_{(1, \sigma^{-1}(r, h))}^*$ or $x_r = c_{(s, \sigma^{-1}(r, h))}^*$. We divide the proof in two steps

- i) The R-parity of the subsequence preceding I_r of the h -symbolic sequence is assumed to be even. Therefore, the R-parity of the subsequence preceding to I_r of the hs -symbolic sequence is also even. Then it follows by using lemma 2, that the entry order from $H_{\sigma^{-1}(1, h)}^*$ to $H_{\sigma^{-1}(r, h)}^*$ is conserved, as $x_1 = c_{(1, 1)}^*$ it results $x_r = c_{(1, \sigma^{-1}(r, h))}^*$, and applying proposition 4, it yields that $H_{\sigma^{-1}(r, h)}^*$ is a convex box. Given that $f^m(C) = f^{nh}(f^r(C)) = f^{nh}(x_r)$ it results that we have moved n times inside the convex box $H_{\sigma^{-1}(r, h)}^*$ from $x_r = c_{(1, \sigma^{-1}(r, h))}^*$. After n movements from $c_{(1, \sigma^{-1}(r, h))}^*$ in a convex box we will be located at $c_{(\sigma^{-1}(n+1, s), \sigma^{-1}(r, h))}^*$ and, using definition 4, it yields $c_{(\sigma^{-1}(n+1, s), \sigma^{-1}(r, h))}^* = C_{((\sigma^{-1}(r, h)-1)s + \sigma^{-1}(n+1, s), hs)}^*$, so we finally obtain

$$\sigma^{-1}(m, hs) = (\sigma^{-1}(r, h) - 1)s + \sigma^{-1}(n + 1, s)$$

- ii) The R-parity of the subsequence preceding I_r of the h -symbolic sequence is assumed to be odd. The proof runs as shown above, the reader should notice that now $x_r = c_{(s, \sigma^{-1}(r, h))}^*$ because the R-parity of the subsequence preceding I_r of the hs -symbolic sequence is odd. The entry order is not conserved and $H_{\sigma^{-1}(r, h)}^*$ is a concave box (by proposition 4). The movements inside $H_{\sigma^{-1}(r, h)}^*$ are ruled by $\overline{\sigma_s^{-1}}$, so after n movements from $c_{(s, \sigma^{-1}(r, h))}^*$ we will be located at $c_{(\overline{\sigma^{-1}}(n+1, s), \sigma^{-1}(r, h))}^*$ and we obtain, by using definition 4,

$$\sigma^{-1}(m, hs) = (\sigma^{-1}(r, h) - 1)s + \overline{\sigma^{-1}}(n + 1, s)$$

Finally taking into account that $\overline{\sigma^{-1}}(n + 1, s) = s + 1 - \sigma^{-1}(n + 1, s)$ (see appendix) statement of the theorem is obtained.

Example 5 We are going to calculate $\sigma_3 \circ \sigma_4$ ($h = 3, s = 4$) to obtain $\sigma_{3 \cdot 4} = \sigma_{12}$. According to figures 1 and 2

$$\sigma_3^{-1} = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix} \quad \sigma_4^{-1} = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 4 & 3 & 2 \end{pmatrix}$$

that give the symbolic sequences RLC and RLLC respectively (see [4] for more details).

- a) If $m = 4$ then $m = 1 \cdot 3 + 1 \equiv n \cdot h + r$. The subsequence preceding $I_r = I_1$ is C, therefore the R-parity is even, hence

$$\sigma^{-1}(4, 12) = (\sigma^{-1}(1, 3) - 1)4 + \sigma^{-1}(2, 4) = 4$$

that is, we are located at $C_{(4,12)}^*$ (see figure 4 and 5.)

- b) If $m = 5$ then $m = 1 \cdot 3 + 2 \equiv n \cdot h + r$. The subsequence preceding $I_r = I_2$ is R, therefore the R-parity is odd, hence

$$\sigma^{-1}(5, 12) = (\sigma^{-1}(2, 3) - 1)4 + 4 + 1 - \sigma^{-1}(2, 4) = 9$$

that is, we are located at $C_{(9,12)}^*$ (see figure 4 and 5), in fact it is just one movement apart from $C_{(4,12)}^*$, obtained when $m = 4$.

Theorem 6 Let the supercycles O_h and O_s have cardinal points $\{C_{(1,h)}^*, \dots, C_{(h,h)}^*\}$ and $\{C_{(1,s)}^*, \dots, C_{(s,s)}^*\}$ respectively. Let σ_h and σ_s be their cardinal ordering permutations. Then the number of iterates to reach $C_{(i,hs)}^*$, with $i = ns + r$, $1 \leq i \leq hs$, and $1 \leq r \leq s$ is given by,

- i) if the R-parity of the subsequence preceding I_{n+1} of the h -symbolic sequence is even,

$$\sigma(i, hs) = (\sigma(r, s) - 1)h + \sigma(n + 1, h)$$

- ii) if the R-parity of the subsequence preceding I_{n+1} of the h -symbolic sequence is odd,

$$\sigma(i, hs) = (\overline{\sigma(r, s)} - 1)h + \sigma(n + 1, h)$$

Proof 6 Let us suppose that the R-parity of the h -sequence preceding I_{n+1} is even. From definition 4

$$C_{(i,hs)}^* = C_{(ns+r,hs)}^* = c_{(r,n+1)}^*$$

Therefore $C_{(i,h)}^*$ is in the r -th position of H_{n+1}^* .

On one hand, $\sigma(n + 1, h)$ iterates are needed, starting from C, to reach the box H_{n+1}^* , furthermore we will be spotted in $c_{(1,n+1)}^* \in H_{n+1}^*$, according lemma 2.

On the other hand, we need $\sigma(r, s) - 1$ iterates inside the box H_{n+1}^* to move from $c_{(1,n+1)}^*$ to $c_{(r,n+1)}^*$ (by definition of $\sigma(r, s)$ and of $c_{(-,-)}^*$, see the paragraph following definition 4). But to move from $c_{(j,n+1)}^*$ to $c_{(j+1,n+1)}^*$ the h boxes need to be visited before; therefore, we need $h(\sigma(r, s) - 1)$ iterates to move from $c_{(1,n+1)}^*$ to $c_{(r,n+1)}^*$.

Finally, we obtain that the total number of iterates, to reach $c_{(r,n+1)}^*$ from C is

$$\sigma(i, hs) = (\sigma(r, s) - 1)h + \sigma(n + 1, h)$$

Conversely if we suppose that the R -parity of the h -sequence preceding I_{n+1} is odd, then the proof can be handled in much the same way, being the only difference that the box H_{n+1}^* is a concave one. We know that H_{n+1}^* is a concave box by using the R -parity of the subsequence preceding I_{n+1} of the h -symbolic sequence. So we have to use $\overline{\sigma(r, s)}$ instead of $\sigma(r, s)$ to control movements inside the box, therefore it results

$$\sigma(i, hs) = (\overline{\sigma(r, s)} - 1)h + \sigma(n + 1, h)$$

The relation between $\sigma(r, s)$ and $\overline{\sigma(r, s)}$ can be seen in the appendix.

Example 6 How many iterates are needed to reach $C_{(9,3 \cdot 4)}^*$? According to example 5, five iterates are needed. Let us check it: if $i = 9 = 2 \cdot 4 + 1 \equiv n \cdot s + r$ then the R -parity of the subsequence preceding $I_{n+1} = I_3$ of the 3-symbolic sequence is odd, hence

$$\sigma(9, 12) = (\overline{\sigma(1, 4)} - 1)3 + \sigma(3, 3) = (\sigma(4, 4) - 1)3 + \sigma(3, 3) = 5$$

as it was expected.

Needless to say in the theorems 5 and 6 we have only considered an iteration number less than or equal to hs , because full cycles are despised.

4 Algorithms

Theorems 5 and 6 can be easily written in an algorithmic manner. The goal is direct when we transform theorem 5 into an algorithm, because as $m = n \cdot h + r$, with $1 \leq r \leq h$, if m is increased one at a time then n keeps constant for h iterates and only r varies, therefore $\sigma^{-1}(n + 1, s)$ keeps constant for h times and only $\sigma^{-1}(r, h)$ varies. The same works in a similar way for theorem 6 with $i = n \cdot s + r$ $1 \leq r \leq s$, where we work with $\sigma(n + 1, h)$ and $\sigma(r, s)$.

4.1 Algorithm to obtain σ_{hs}^{-1} from σ_h^{-1} and σ_s^{-1}

- (1) Write the h -symbolic sequence.
- (2) Write $\sigma^{-1}(j, s)$ below every symbol of the h -symbolic sequence.
- (3) Change $\sigma^{-1}(j, s)$ by $s + 1 - \sigma^{-1}(j, s)$ if the R -parity of the previous sequence of upper symbols is odd.

- (4) Add $(\sigma^{-1}(1, h) - 1) \cdot s$ to the first term, $(\sigma^{-1}(2, h) - 1) \cdot s$ to the second term and so successively until you add $(\sigma^{-1}(h, h) - 1) \cdot s$ to the last term.
- (5) Repeat consecutively the former sequence “ s ” times, with “ j ” changing from $j = 1$ to $j = s$.

The sequence obtained in step five is the looked for permutation.

Example 7 We are going to obtain $\sigma_{3,4}^{-1}$ from σ_3^{-1} and σ_4^{-1}

(1) R L C

(2)

$$\begin{array}{ccc} \text{R} & \text{L} & \text{C} \\ \sigma^{-1}(j, 4) & \sigma^{-1}(j, 4) & \sigma^{-1}(j, 4) \end{array}$$

(3)

$$\sigma^{-1}(j, 4) \quad s + 1 - \sigma^{-1}(j, 4) \quad s + 1 - \sigma^{-1}(j, 4) \implies \sigma^{-1}(j, 4) \quad 5 - \sigma^{-1}(j, 4) \quad 5 - \sigma^{-1}(j, 4)$$

(4)

$$\begin{array}{ccc} \sigma^{-1}(j, 4) & 5 - \sigma^{-1}(j, 4) & 5 - \sigma^{-1}(j, 4) \\ + & + & + \\ (\sigma^{-1}(1, 3) - 1) \cdot 4 & (\sigma^{-1}(2, 3) - 1) \cdot 4 & (\sigma^{-1}(3, 3) - 1) \cdot 4 \\ \implies \sigma^{-1}(j, 4) & 13 - \sigma^{-1}(j, 4) & 9 - \sigma^{-1}(j, 4) \end{array}$$

(5)

$$\begin{array}{c} \sigma^{-1}(1, 4) \quad 13 - \sigma^{-1}(1, 4) \quad 9 - \sigma^{-1}(1, 4) \quad \left| \sigma^{-1}(2, 4) \quad 13 - \sigma^{-1}(2, 4) \quad 9 - \sigma^{-1}(2, 4) \right| \\ \left| \sigma^{-1}(3, 4) \quad 13 - \sigma^{-1}(3, 4) \quad 9 - \sigma^{-1}(3, 4) \right| \quad \left| \sigma^{-1}(4, 4) \quad 13 - \sigma^{-1}(4, 4) \quad 9 - \sigma^{-1}(4, 4) \right| \implies \\ \implies 1 \quad 12 \quad 8 \quad 4 \quad 9 \quad 5 \quad 3 \quad 10 \quad 6 \quad 2 \quad 11 \quad 7 \end{array}$$

It can be checked in figures 4 and 5 that we have obtained $\sigma_{3,4}^{-1}$ (the visiting sequence of $O_{3,4}$)

4.2 Algorithm to obtain σ_{hs} from σ_h and σ_s

- (1) Write “ s ” times $\sigma(j, h)$.
- (2) Add $(\sigma(1, s) - 1) \cdot h$ to the first term, $(\sigma(2, s) - 1) \cdot h$ to the second term and so successively until you add $(\sigma(s, s) - 1) \cdot h$ to the last term
- (3) Repeat consecutively the former sequence “ h ” times, with “ j ” changing from $j = 1$ to $j = h$.

- (4) If the R -parity of the subsequence preceding $\underline{I_j}$ of the h -symbolic sequence is odd then $\sigma(i, s)$ must change by $\underline{\sigma(i, s)}$ for $i = 1, \dots, s$.

The obtained sequence is the looked for permutation.

Example 8 *We are going to obtain $\sigma_{3,4}$ from σ_3 and σ_4*

$$(1) \quad \sigma(j, 3) \quad \sigma(j, 3) \quad \sigma(j, 3) \quad \sigma(j, 3)$$

(2)

$$\begin{array}{cccc} \sigma(j, 3) & \sigma(j, 3) & \sigma(j, 3) & \sigma(j, 3) \\ + & + & + & + \\ (\sigma(1, 4) - 1) \cdot 3 & (\sigma(2, 4) - 1) \cdot 3 & (\sigma(3, 4) - 1) \cdot 3 & (\sigma(4, 4) - 1) \cdot 3 \end{array}$$

(3)

$$\begin{array}{cccc} \sigma(1, 3) & \sigma(1, 3) & \sigma(1, 3) & \sigma(1, 3) \\ + & + & + & + \\ (\sigma(1, 4) - 1) \cdot 3 & (\sigma(2, 4) - 1) \cdot 3 & (\sigma(3, 4) - 1) \cdot 3 & (\sigma(4, 4) - 1) \cdot 3 \end{array}$$

$$\begin{array}{cccc} \sigma(2, 3) & \sigma(2, 3) & \sigma(2, 3) & \sigma(2, 3) \\ + & + & + & + \\ (\sigma(1, 4) - 1) \cdot 3 & (\sigma(2, 4) - 1) \cdot 3 & (\sigma(3, 4) - 1) \cdot 3 & (\sigma(4, 4) - 1) \cdot 3 \end{array}$$

$$\begin{array}{cccc} \sigma(3, 3) & \sigma(3, 3) & \sigma(3, 3) & \sigma(3, 3) \\ + & + & + & + \\ (\sigma(1, 4) - 1) \cdot 3 & (\sigma(2, 4) - 1) \cdot 3 & (\sigma(3, 4) - 1) \cdot 3 & (\sigma(4, 4) - 1) \cdot 3 \end{array}$$

(4)

$$\begin{array}{cccc}
\begin{array}{c} \sigma(1,3) \\ + \\ (\sigma(1,4) - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(1,3) \\ + \\ (\sigma(2,4) - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(1,3) \\ + \\ (\sigma(3,4) - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(1,3) \\ + \\ (\sigma(4,4) - 1) \cdot 3 \end{array} \\
\\
\begin{array}{c} \sigma(2,3) \\ + \\ (\overline{\sigma(1,4)} - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(2,3) \\ + \\ (\overline{\sigma(2,4)} - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(2,3) \\ + \\ (\overline{\sigma(3,4)} - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(2,3) \\ + \\ (\overline{\sigma(4,4)} - 1) \cdot 3 \end{array} \\
\\
\begin{array}{c} \sigma(3,3) \\ + \\ (\overline{\sigma(1,4)} - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(3,3) \\ + \\ (\overline{\sigma(2,4)} - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(3,3) \\ + \\ (\overline{\sigma(3,4)} - 1) \cdot 3 \end{array} &
\begin{array}{c} \sigma(3,3) \\ + \\ (\overline{\sigma(4,4)} - 1) \cdot 3 \end{array} \\
\Rightarrow 1 & 10 & 7 & 4 & 6 & 9 & 12 & 3 & 5 & 8 & 11 & 12
\end{array}$$

It can be checked that the obtained permutation $\sigma_{3.4}$ is the inverse permutation of $\sigma_{3.4}^{-1}$ obtained in example 7. Observe that the added sequence $\sigma(i, s)$ is always the same or a mere reflection of it (see appendix) so the algorithm is very fast to run.

5 Appendix

5.1 Meaning of the conjugation for permutations σ_p and σ_p^{-1}

Let us consider two unimodal functions, with a maximum and a minimum respectively, such that they have the same orbits. It is well known that their symbolic sequences are conjugated by the change $R \longleftrightarrow L$. The conjugation can be seen as a homotecy that transforms the maxima into a minima and viceversa. If we pay attention to figure 6 we can see as the sequence RLC (in the maximum) is transform into LRC (in the minimum). We can interprete geometrically the homotecy as a reflection about the dashed line. We want to know how the reflection affects to σ_p and σ_p^{-1} .




Figure 6. Orbit, with symbolic sequence RLC, in the first quadrant is transformed into an orbit with symbolic sequence LRC in the third quadrant by means of a homotety. The homotety is performed as a reflection about the dashed line.

- The permutation

$$\sigma_p = \begin{pmatrix} 1 & \cdots & i & \cdots & p \\ \sigma(1, p) & \cdots & \sigma(i, p) & \cdots & \sigma(p, p) \end{pmatrix} \quad (2)$$

is defined by $f^{\sigma(i,p)} = C_i^*$ (see [4]) therefore, its reflection is given by

$$\overline{\sigma}_p = \begin{pmatrix} 1 & \cdots & i & \cdots & p \\ \sigma(p, p) & \cdots & \sigma(p-i+1, p) & \cdots & \sigma(1, p) \end{pmatrix} \quad (3)$$

that is,

$$\begin{pmatrix} 0 & \cdots & 0 & 1 \\ 0 & \cdots & 1 & 0 \\ \vdots & \ddots & \vdots & 0 \\ 1 & \cdots & \cdots & 0 \end{pmatrix} \begin{pmatrix} \sigma(1, p) \\ \sigma(2, p) \\ \vdots \\ \sigma(p, p) \end{pmatrix} = \begin{pmatrix} \sigma(p, p) \\ \sigma(p-1, p) \\ \vdots \\ \sigma(1, p) \end{pmatrix}$$

where the matrix $\begin{pmatrix} 0 & \cdots & 0 & 1 \\ 0 & \cdots & 1 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 1 & \cdots & \cdots & 0 \end{pmatrix}$ accomplishes the reflection.

- The permutation σ_p^{-1} determines how the p -periodic orbit is visited (see [4]), that is, what cardinal points are visited one after another. The reflection of σ_p^{-1} is $\overline{\sigma}_p^{-1}$ that means the orbit is gone through in reverse order.

Therefore one of the permutations produces a shift which is cancelled by the other, hence

$$\sigma^{-1}(i, p) + \overline{\sigma^{-1}(i, p)} = p + 1$$

σ_p^{-1} and $\overline{\sigma_p^{-1}}$ are deduced from (2) and (3)

6 Conclusions

To understand the language of chaos we need to understand its grammar. A grammar, a set of rules, that determines how to combine some structures with others to generate new ones. In nonlinear dynamics basic structures are orbits, that are characterized by their cardinal ordering permutations (COP) [4]; whereas, grammar is the COP composition rules.

These COP composition rules, unknown until now, have been stated in this paper in two theorems. Such composition rules have two important and immediate implications:

- (i) Given the COP of a h -periodic and s -periodic orbit then the COP of hs -periodic orbit is deduced, that is, the relative position of all points belonging to the hs -periodic orbit and its visiting order. Furthermore using the composition law in a recursive way it is possible to obtain COP for $h^a s^b$ -periodic orbits $a, b \in \mathbb{Z}^+$, that is, with a couple of COP we can determine infinitely many COP and from them to fully understand the associated orbits.
- (ii) As COP for periodic doubling cascade is known [4] we would immediately obtain COP for both periodic doubling cascade and saddle-node bifurcation cascade [9,10] with any arbitrary basic period $(p \cdot 2^k)$.

Known the points (i) and (ii) all underlying periodic structures in an unimodal function f are determined, hence the importance of the theorems 5 and 6.

We want to point out that the COP for the period doubling cascade can be obtained by using the theorems proven in this paper. It is enough considering $\sigma_h = \sigma_s$ ($h = s = 2$) to obtain $\sigma_{h.s} = \sigma_{2.2}$. By using recursively this process we get the COP for the 2^n -periodic orbit of the cascade. No matter whether we take $h = 2^{(n-1)}$ and $s = 2$ or $h = 2$ and $s = 2^{(n-1)}$; in this case the process is absolutely symmetric. But if we use the second one we get a new proof of theorem 1 in [4].

To prove the theorems we needed to know if some points of h -periodic orbit were associated with maxima or minima of f^h . Since it would be wholly inoperative to get that information by deriving f^h when h is big, two steps

were carried on: a) to provide this information implicitly in the theorems, b) to obtain this information in an simple way, even if h is very big (proposition 4). The solutions to these problems play a pivotal role in proving those theorems. We must point out that analytical information has been obtained from topological one (what authors named “subsequence preceding”), establishing a link between two apparently unconnected fields, that must be studied more carefully. It is well known that, when two apparently unconnected mathematical fields turn out to describe the same thing, but from two different points of view, the problems they address are very largely simplified.

References

- [1] HAO BAI-LIN : *Elementary Symbolic dynamicss*. World Scientific 1989.
- [2] N. METROPLOIS, M. L. STEIN, AND P. R. STEIN: *On Finite Limit Sets for Transformations on the Unit Interval*, J. Combinatorial Theory **15** (1973) 25-44
- [3] B. DERRIDA, A. GERVOIS, Y. POMEAU : *Iteration of endomorphisms on the rael axes and representation of numbers*, L’I.H.P. section A **29** 3, (1978) 305-356
- [4] J. SAN MARTÍN, M^A JOSÉ MOSCOSO, A. GONZÁLEZ GÓMEZ: *The universal cardinal ordering of fixed points*, Chaos, Solitons & Fractals, **42** (2009) 1996-2007
- [5] M. J. FEIGENBAUM: *Quantitative Universality for a Class of Nonlinear Transformations*, J. Stat. Phys. **19** (1978) 25-52
- [6] M. J. FEIGENBAUM: *The Universality Metric Properties for Nonlinear Transformations*, J. Stat. Phys. **21** (1979) 669-706
- [7] J. MILNOR, W. THURSTON: *On iterated maps of the interval*, Dinamical systems. Lecture notes in math, vol. 1342. Berlin: Springer; (1988) 465-563.
- [8] R. GILMOR, M. LEFRANC: *The topology of chaos*, New York: Willye; (2002).
- [9] J. SAN MARTÍN, D. RODRÍGUEZ-PÉREZ: *Intermittency cascade*, Chaos, Solitons & Fractals, **32**(2) (2007) 816-831
- [10] J. SAN MARTÍN, D. RODRÍGUEZ-PÉREZ: *Conjugation of cascades*, Chaos, Solitons & Fractals, **39**(2) (2009) 666-681
- [11] GLEN R. HALL: *Some examples of permutations modellings area preserving monotone twist map*, Physica D: Nonlinear Phenomena, **28** (1978) 393-400
- [12] J. M. AMIGÓ, M. B. KENNEL, L. KOCAREV : *The permutations entropy rate equals the metric entropy rate for ergodic information sources and ergodic dynamical systems*, Physica D **210**(2005) 77-95.

- [13] J. M. AMIGÓ, L. KOCAREV, I. TOMOVSKI : *Discrete entropy* , 228(2007) 77-85
- [14] J. M. AMIGÓ, M. B. KENNEL: *Topological permutations entropy* , Physica D 231(2007) 137-142.
- [15] C. BANDT, G. KELLER, B. POMPE : *Entropy of interval maps via permutations*, Nonlinearity 15(2002) 1595-1602
- [16] J. M. AMIGÓ, M. B. KENNEL: *Forbidden ordinal patterns in higher dimensional dynamics*, Physica D 237(2008) 2893-2899.
- [17] J. M. AMIGÓ, L. KOCAREV, J. SZCZEPANSKI : *Order pattern and chaos*, Physics Letters A 355(2006) 27-31.
- [18] L. ZUNINO, M. ZANIN, B. M. TABAK, D.G. PÉREZ: *Forbidden patterns, permutation entropy and stock market inefficiency*, Physica A 388(2009) 2854-2864.